

Alberta-Thy-09-99
hep-th/9910158

Comments on “ Black Hole Entropy from Conformal Field Theory in Any Dimension ”

by

Mu-In Park^{a1} and Jeongwon Ho^b

^a *Department of Physics, Yonsei University, Seoul 120-749, Korea*

^b *Theoretical Physics Institute, Department of Physics, University of Alberta, Edmonton,
Canada T6G 2J1*

ABSTRACT

In a recent letter, Carlip proposed a generalization of the Brown-Henneaux-Strominger construction to any dimension. We present two criticisms about his formulation.

September 1999

¹Electronic address: mipark@physics3.sogang.ac.kr

Recently Carlip [1] proposed a generalization of the Brown-Henneaux-Strominger construction to any dimension based on the algebra of deformations at a black hole horizon. By restricting to the horizon of a black hole, he has shown that the algebra contains a Virasoro subalgebra independent of the spatial dimensions with the appropriate boundary conditions. With the aid of Cardy formula he derived the Bekenstein-Hawking entropy within the conformal field theory. In this Comment, we point out that (a) his analysis has a serious error which invalidates one of its principal claims and (b) independent of that error there is one more serious defect.

The error (a) occurs in $\delta_{\xi_2} J[\hat{\xi}_1]$ (10) of Ref. [1] which essentially claims that it should be antisymmetric under $1 \leftrightarrow 2$ interchange according to his identification of $\delta_{\xi_2} J[\hat{\xi}_1] = \{J[\hat{\xi}_2], J[\hat{\xi}_1]\}^*$ ($\{, \}^*$ denotes the Dirac bracket). But this is not correct identification because the antisymmetry can not be attained with his choice of $J[\hat{\xi}]$ (9) and hence the Virasoro algebra, whose existence is one of its principal claims, can not be obtained in his form. Let us explain our conclusion in detail. We first note that the last two terms of $\delta_{\xi_2} J[\hat{\xi}_1]$ have no manifest antisymmetry for indices. In his paper [1], he “assumed” $\delta_{\xi_2} \hat{\xi}_1^t = \hat{\xi}_2^r \partial_r \hat{\xi}_1^t$ to get the antisymmetry “by hand”. But, this assumption is not acceptable because one can show that $\delta_{\xi_2} \hat{\xi}_1^t = 0$ which is essentially because $\delta_{\xi_2} \hat{\xi}_1^t = (\delta_{\xi_2} N) \xi_1^t = 0$. Hence, the $\delta_{\xi_2} J[\hat{\xi}_1]$ is not antisymmetric under $1 \leftrightarrow 2$ because of the last two terms although it is antisymmetric for the first two terms in (13). One can not obtain the Virasoro algebra with this preliminary algebra.

About the origin of this failure, we found that it is a direct consequence of the *non-differentiability* of $L = H + J$ contrast to his claim. δJ does not cancel the boundary terms in δH exactly, rather it introduces several additional boundary terms in δL : Most of these terms have no contribution on the horizon because of his boundary condition $N^2|_{r_+} = 0$ but two of them can not be neglected and these two contributions make L non-differentiable. Let us explain first why the differentiability of L is connected to the antisymmetrization of $\delta_{\xi_2} J[\hat{\xi}_1]$. When L is the correct diffeomorphism (*Diff*) generator, i.e., $\{L[\eta], \phi\} = \delta_\eta \phi$ for any dynamical variables ϕ , the *Diff* transformation of L can be expressed as $\delta_\eta L[\xi] = \{L[\eta], L[\xi]\}$. When the constraint $H \approx 0$ is imposed this will reduce to : $\delta_\eta J[\xi] = \{J[\eta], J[\xi]\}^*$ by picking out the boundary parts of $\delta_\eta L[\xi]$ according to the definition of the Dirac bracket. According to this identification, $\delta_\eta J[\xi]$ is antisymmetric under $\eta \leftrightarrow \xi$ explicitly as $\delta_\eta L[\xi]$ is. Now, let us consider the (general) variation of L to investigate its differentiability:

$$\begin{aligned} \delta L &= \frac{1}{8\pi G} \int_{r_+} d^{n-2} x \sqrt{\sigma} \left[\frac{1}{2} \sigma^{\alpha\beta} n^r \hat{\xi}^t D_r \delta \sigma_{\alpha\beta} - n_r \hat{\xi}^r \delta K + \dots \right] + \text{bulk} \\ &= -\frac{1}{8\pi G} \int d^{n-1} x \delta(r - r_+) \sqrt{\sigma} \left[\frac{\sigma^{\alpha\beta}}{2} \partial_r (n^r \hat{\xi}^t) \delta g_{\alpha\beta} + \frac{n_r \hat{\xi}^r}{(2-n)f\sqrt{\sigma}} g_{rr} \delta \pi^{rr} + \dots \right] + \text{bulk}, \end{aligned}$$

where the contributions of ‘...’ terms to the *Diff* transformation $\delta_\eta \phi$ are order of $O(N^2)$ or $O(N)$ which vanish on the horizon. From these two non-vanishing boundary contributions, we

get the corresponding *anomalous* boundary contributions in the following *Diff* transformations:

$$\begin{aligned}
\delta_\xi g_{rr}(x) &= \{L[\hat{\xi}], g_{rr}(x)\} \\
&= \frac{1}{8\pi G} \delta(r - r_+) \frac{n_r \hat{\xi}^r}{(2 - n)f\sqrt{\sigma}} g_{rr} + \text{bulk}, \\
\delta_\xi \pi^{\alpha\beta}(x) &= \{L[\hat{\xi}], \pi^{\alpha\beta}(x)\} \\
&= -\frac{1}{8\pi G} \delta(r - r_+) \frac{\sqrt{\sigma}}{2} \sigma^{\alpha\beta} (n^r \hat{\xi}^t) + \text{bulk},
\end{aligned}$$

whose boundary contributions behave as $\delta(r - r_+) \times O(1)$ which can not be neglected. Here, the bulk part will represent the usual *Diff* transformations in the bulk. These wrong *Diff* transformations implies that Carlip's original choice of $J[\hat{\xi}]$ is not correct one to make $L[\hat{\xi}]$ be the *Diff* generator for the system with the horizon boundary and to give the claimed Virasoro algebra.

Criticism (b) consists of the observation that even upon ignoring point (a), his scenario does not work for the non-rotating black hole even though it works well for the rotating case. Recently a resolution has been suggested in a different context of 2D gravity by Cadoni and Mignemi [2] who have introduced the time-integrated generator instead of (usual) fixed-time generator. But we do not think that this can be resolved even in the way of Cadoni and Mignemi although the time integration of (21) for the non-rotating case gives the similar result as the rotating case with different central charge $c = 3A\beta/(\pi GT)$ and different $L_0 = AT/(8\pi G\beta)$ [3]. A strange fact is that the resulting entropy $\log\rho(L_0) = 2 \times \frac{A}{4G}$ has a wrong factor of 2 which remind us a similar wrong factor in their computation of 2D-gravity entropy [2]. More serious problem of this method is that the time integration of the left-hand sides of (21) can not be written as $\{\bar{J}[\hat{\xi}_m], \bar{J}[\hat{\xi}_n]\}^*$ for time-integrated generator $\bar{J} = \frac{1}{\tau} \int_0^\tau dt J(t)$, which is essential for the interpretation of (21) as a Virasoro algebra.

MIP would like to thank Prof. Steven Carlip for stimulus discussions along this work and Prof. Jae Hyung Yee, Drs. Gung-won Kang and Hyuk-jae Lee for several helpful discussions. MIP was supported in part by a postdoctoral grant from the Natural Science Research Institute, Yonsei University in the year 1999 and in part by the Korea Research Foundation under 98-015-D00061. JH was supported in part by Korea Science and Engineering Foundation and in part by National Science and Engineering Research Council of Canada.

References

- [1] S. Carlip, Phys. Rev. Lett. **82**, 2828 (1999).
- [2] M. Cadoni and S. Mignemi, Phys. Rev. **D59**, 081501 (1999).

[3] M.-I. Park and J. H. Yee (unpublished).